

Adaptions of oxidative phosphorylation to acute and chronic exercise

Alvarez-Luis J¹, Khatbane K¹, Gama-Perez P¹, Cutanda-Tesouro S¹, Cardoso LHD², Komlódi T³, Bosch N¹, Barrientos R¹, Leon B¹, Neri L¹, Sivasailam A⁴, Fernandez-Montes P⁵, Irving B⁴, Enriquez JA⁵, Gnaiger E², Garcia-Roves Pablo M¹

¹Dept. Physiological Sciences, Univ. of Barcelona, and Bellvitge Biomedical Research Institute (IDIBELL) Barcelona, Spain; ²Oroboros Instruments, Innsbruck, Austria; ³Dept. of Biochemistry, Semmelweis Univ., Budapest, Hungary; ⁴Dept. of Kinesiology, Univ. of Virginia, Virginia, USA; ⁵Spanish National Center for Cardiovascular Research, Madrid, Spain. pgarciaroves@ub.edu

Contribution to [MiP2026](#) – Received 2026-09-28 – Online 2026-10-07

Cite: Alvarez-Luis J, Khatbane K, Gama-Perez P, Cutanda-Tesouro S, Cardoso LHD, Komlódi T, Bosch N, Barrientos R, Leon B, Neri L, Sivasailam A, Fernandez-Montes P, Irving B, Enriquez JA, Gnaiger E, Garcia-Roves Pablo M (2026) Adaptions of oxidative phosphorylation to acute and chronic exercise. BEC preprints 2026.10 <https://doi.org/10.26124/becprep2026-0010>

Introduction

Skeletal muscle is a highly adaptable tissue that continuously remodels in response to physiological stimuli, particularly exercise-induced contractile activity. Because muscle contraction imposes substantial energetic demands, mitochondrial function and oxidative phosphorylation (OXPHOS) play central roles in maintaining ATP production, metabolic flexibility, and overall muscle performance. Physiological mitochondrial adaptations, including improved respiratory capacity, are key mechanisms underlying enhanced exercise performance (Garcia-Roves et al 2025), whereas mitochondrial dysfunction is associated with aging, obesity, and metabolic disease (Georgiev et al 2022).

High-resolution respirometry (HRR) provides sensitive and direct measurements of oxygen consumption. The use of permeabilized muscle fibers (pfi) offers advantages compared to isolated mitochondria, including preservation of mitochondrial architecture and maintenance of both subsarcolemmal and intermyofibrillar mitochondrial populations. Recent technological advances expand mitochondrial phenotyping beyond respiration, enabling simultaneous measurements of mitochondrial membrane potential ($\Delta\psi_{mt}$), ATP production, reactive oxygen species generation, and coenzyme Q (CoQ) redox state.

The Oroboros NextGen-O2k Q-Module enables real-time monitoring of both oxygen consumption and CoQ redox status. As CoQ acts as the central electron carrier linking Complexes I and II to Complex III, its redox state provides valuable information about electron transport system activity and mitochondrial metabolic status. This approach has been successfully applied in isolated mitochondria (Komlódi et al. 2021). The

feasibility of combining HRR, CoQ redox measurements, and membrane potential assessment in skeletal muscle pfi was a challenge faced in the present study.

First, we optimized an integrated platform for the simultaneous determination of mitochondrial respiratory capacity, CoQ redox state, and membrane potential in pfi. Second, applying this platform we investigated mitochondrial adaptations induced by acute and chronic exercise. This approach provides new insights into the mechanisms underlying exercise-induced improvements in mitochondrial function.

Methods

Experiments were conducted on young adult male C57BL/6J mice maintained under standardized laboratory conditions. Extensor digitorum longus (EDL) muscles were excised, mechanically separated into fiber bundles, and chemically permeabilized with saponin. This procedure provided direct access to the mitochondrial network while preserving muscle structure and mitochondrial organization (Doerrier et al 2024).

Mice were assigned to three experimental groups: sedentary control, acute exercise, and chronic exercise training. The chronic group underwent a 4-week training protocol (Gonzalez-Franquesa et al 2022). The acute exercise protocol consisted of treadmill running at a 5° incline for 10 min at $0.17 \text{ m}\cdot\text{s}^{-1}$, followed by 10 min at $0.2 \text{ m}\cdot\text{s}^{-1}$ and 10 min at $0.23 \text{ m}\cdot\text{s}^{-1}$. Thereafter, animals performed repeated 2-min intervals at progressively increasing speeds (0.27, 0.3, and $0.33 \text{ m}\cdot\text{s}^{-1}$, each repeated twice), interspersed with 2-min recovery periods at $0.2 \text{ m}\cdot\text{s}^{-1}$.

Mitochondrial respiration was assessed using the Oroboros NextGen-O2k with the substrate-uncoupler-inhibitor titration protocol SUIT-008 (https://www.bioblast.at/index.php/SUIT-008_O2_pfi_D014), allowing sequential evaluation of leak respiration, OXPHOS capacity, electron transfer (ET) capacity, and residual oxygen consumption. Oxygen consumption rates were corrected for instrumental background and residual oxygen consumption and normalized to tissue mass.

The redox state of the mitochondrial CoQ pool was monitored using the NextGen-O2k Q-Module and a short-chain CoQ analogue (CoQ_2). The oxidation and reduction peak potentials were determined by cyclic voltammetry at $+70 \pm 12 \text{ mV}$ and $-259 \pm 10 \text{ mV}$, respectively. Changes in electrical current generated by oxidation of reduced CoQ_2 are indirect indicators of CoQ redox status (Komlódi et al 2021).

Mitochondrial membrane potential ($\Delta\psi_{\text{mt}}$) was measured by safranin fluorescence coupled to the O2k Fluo-Module. Fluorescence changes across respiratory states were used to evaluate relative changes in membrane potential (Krumschnabel et al 2014).

Mitochondrial protein content and markers of mitochondrial dynamics were analyzed by SDS-PAGE and immunoblotting using standard procedures. To evaluate respiratory complexes and supercomplex organization, blue native electrophoresis and in-gel

activity assays were performed in isolated mitochondrial preparations (Acín-Pérez et al 2020).

Data were analyzed using DatLab 8 and GraphPad Prism. Statistical comparisons were performed using non-parametric tests when appropriate.

Results

Optimization of simultaneous measurements: Previously reported CoQ₂ concentrations interfered with respiration in pfi. Therefore, a concentration-response analysis was performed. Concentrations ≥ 40 nM significantly inhibited oxygen consumption, whereas 20 nM CoQ₂ exerted no detectable effect. Validation using the SUIT-008 protocol confirmed that 20 nM CoQ₂ did not alter leak respiration, OXPHOS, or ET capacity and was selected for subsequent experiments.

The suitability of 1 μ M safranin for membrane potential measurements was evaluated. Safranin produced robust and reproducible fluorescence changes across respiratory states, confirming adequate sensitivity for $\Delta\psi_{mt}$ determination. Importantly, safranin did not significantly affect oxygen flux under most conditions, although a slight inhibitory effect was observed on NADH-linked OXPHOS respiration, consistent with previous reports (Krumschnabel et al 2014).

Together, these results established a platform for simultaneous quantification of mitochondrial respiration and either CoQ redox state or mt-membrane potential in pfi.

Effects of exercise on mitochondrial respiration: Mitochondrial bioenergetics was compared among sedentary controls, acutely exercised mice, and chronically trained mice. Both acute and chronic exercise significantly increased mitochondrial respiratory capacity. The enhancement was particularly evident in OXPHOS respiration when NADH- and succinate-linked pathways were simultaneously activated (Control: 138.7 ± 9.67 ; Acute: 204.09 ± 28.10 ; Chronic: 195.95 ± 13.80 pmol·s⁻¹·mg⁻¹) and in ET capacity upon uncoupler titrations (Control: 132.1 ± 8.63 ; Acute: 207.67 ± 22.66 ; Chronic: 190.06 ± 15.08 pmol·s⁻¹·mg⁻¹).

Interestingly, the relative contribution of NADH-linked respiration to maximal oxygen flux was reduced significantly after acute but not chronic exercise (Control: 0.759 ± 0.019 ; Acute: 0.635 ± 0.014 ; Chronic: 0.705 ± 0.030 pmol·s⁻¹·mg⁻¹), suggesting exercise-induced remodelling of the ETS.

Effects of exercise on mitochondrial membrane potential: Mitochondrial membrane potential displayed the expected pattern across respiratory states, reaching the highest values during leak respiration and the lowest in fully decoupled conditions. Despite exercise-induced increases in respiratory capacity, neither acute nor chronic exercise significantly altered $\Delta\psi_{mt}$ in any respiratory state. These findings indicate that mt-membrane potential remains tightly regulated despite substantial changes in oxidative capacity.

Effects of exercise on CoQ redox state: Analysis of CoQ redox status revealed transient exercise-induced alterations. Acute exercise made the CoQ junction more oxidized during states selected OXPHOS (NADH and SDH-linked respiration - Control: 0.764 ± 0.026 ; Acute: 0.668 ± 0.023 reduced Q fraction) and ET states (reduced Q fraction of NADH and SDH-linked respiration - Control: 0.796 ± 0.023 ; Acute: 0.702 ± 0.023 ; and reduced Q fraction of SDH-linked respiration - Control: 0.835 ± 0.020 ; Acute: 0.733 ± 0.021), indicating temporary changes in electron flow through the respiratory system. This was largely associated with an improved capacity of succinate-derived electron input into the Q-junction under both substrate-saturated (reduced Q fraction OXPHOS SDH-linked respiration - Control: 0.972 ± 0.033 ; Acute: 0.763 ± 0.040) and submaximal (reduced Q fraction of Leak SDH-linked respiration - Control: 0.941 ± 0.039 ; Acute: 0.721 ± 0.042) conditions. However, these differences were no longer evident after chronic training, with CoQ redox values comparable to controls.

The improvements in respiratory capacity may involve post-translational mechanisms. Protein levels of the mitochondrial dynamics markers MFN2 and DRP1 were unchanged among groups. However, muscles from both acute and chronic exercise groups exhibited increased concentrations of I+III₂ supercomplexes compared with controls, suggesting that OXPHOS reorganization contributes to the observed improvements in mitochondrial bioenergetic performance.

Overall, these findings suggest that acute exercise induces a transient redistribution of redox balance within the ETS, whereas long-term adaptations, including increased mitochondrial content, restore redox homeostasis.

Discussion

The present study demonstrates the successful implementation of a novel experimental approach integrating HRR with simultaneous assessment of mitochondrial membrane potential and CoQ redox state in skeletal muscle pfi. Compared with respiration measurements alone, this methodology provides a more comprehensive evaluation of mitochondrial fitness.

The improvements in mitochondrial respiratory function following chronic exercise are consistent with extensive evidence showing enhanced oxidative capacity after endurance training (Garcia-Roves et al 2025). Increased OXPHOS and ET capacities reflect enhanced mitochondrial performance and support the concept that exercise promotes adaptations that improve aerobic metabolism.

A particularly notable finding is the increase in respiratory capacity after a single exercise bout. Although mitochondrial biogenesis generally requires repeated exercise stimuli, acute improvements in mitochondrial function may arise through more rapid mechanisms, including post-translational regulation, enhanced respiratory

supercomplex assembly, altered mitochondrial dynamics, or increased efficiency of existing respiratory complexes (Garcia-Roves et al 2025).

The analysis of CoQ redox state provides additional insights into these adaptive processes. CoQ serves as a key junction of electron transfer and its analysis allows to infer which part of the ETS is affected. The transient decrease in the fraction of reduced CoQ observed after acute exercise may reflect improved electron transfer efficiency downstream of CoQ or acute reorganization of OXPHOS activity. The restoration of control-like redox values following chronic training suggests the establishment of a new metabolic equilibrium associated with the well-described increase in mitochondrial content.

In contrast, mitochondrial membrane potential remained remarkably stable across all groups. Despite substantial improvements in OXPHOS and ET capacities, $\Delta\psi_{mt}$ was maintained throughout all respiratory states. This finding suggests that exercise-induced enhancements in respiration occur without major alterations in mt-membrane potential and that mitochondrial energy transformation remains tightly regulated.

Collectively, these findings support the concept that exercise enhances mitochondrial respiratory capacity through both quantitative and qualitative improvements in electron transfer efficiency rather than through persistent changes in membrane potential or CoQ redox balance. These adaptations appear to depend on mitochondrial biogenesis and OXPHOS reorganization into more efficient configurations.

Few studies ever examined acute exercise-induced changes in CoQ redox state and membrane potential in skeletal muscle. The present work demonstrates that simultaneous measurement of respiration, membrane potential, and CoQ redox status in pfi is both feasible and reliable. The results show that acute and chronic exercise enhance mitochondrial respiratory capacity, with evidence that even a single exercise session can improve skeletal muscle bioenergetics. While membrane potential remains stable, acute exercise transiently alters CoQ redox balance before normalization with chronic training. Overall, these findings provide novel mechanistic insights into exercise-induced mitochondrial adaptations and establish an integrated methodological framework for future studies of mitochondrial physiology in health and disease.

References

- Acin-Perez R, Hernansanz-Agustin P, Enriquez JA (2020) Analyzing electron transport chain supercomplexes. *Methods Cell Biol* 155:181-97. <https://doi.org/10.1016/bs.mcb.2019.12.002>
- Doerrier C, Gama-Perez P, Pesta D, Distefano G, Soendergaard SD, Maise Chroéis K, Gonzalez-Franquesa A, Goodpaster BH, Prats C, Sales-Pardo M, Guimera R, Coen PM, Gnaiger E, Larsen S, Garcia-Roves PM (2024) Harmonization of experimental procedures to assess mitochondrial respiration in human permeabilized skeletal muscle fibers. *Free Radic Biol Med* 223:384-97. <https://doi.org/10.1016/j.freeradbiomed.2024.07.039>

- Garcia-Roves PM, Alvarez-Luis J, Cutanda-Tesouro S (2025) The role of skeletal muscle respiratory capacity in exercise performance. *Free Radic Biol Med* 229:474-84. <https://doi.org/10.1016/j.freeradbiomed.2024.12.060>
- Georgiev A, Granata C, Roden M (2022) The role of mitochondria in the pathophysiology and treatment of common metabolic diseases in humans. *Am J Physiol Cell Physiol* 322:C1248-59. <https://doi.org/10.1152/ajpcell.00035.2022>
- Gonzalez-Franquesa A, Gama-Perez P, Kulis M, Szczepanowska K, Dahdah N, Moreno-Gomez S, Latorre-Pellicer A, Fernández-Ruiz R, Aguilar-Mogas A, Hoffman A, Monelli E, Samino S, Miró-Blanch J, Oemer G, Duran X, Sanchez-Reboredo E, Schneeberger M, Obach M, Montane J, Castellano G, Chapaprieta V, Sun W, Navarro L, Prieto I, Castaño C, Novials A, Gomis R, Monsalve M, Claret M, Graupera M, Soria G, Wolfrum C, Vendrell J, Fernández-Veledo S, Enríquez JA, Carracedo A, Perales JC, Nogueiras R, Herrero L, Trifunovic A, Keller MA, Yanes O, Sales-Pardo M, Guimerà R, Blüher M, Martín-Subero JI, Garcia-Roves PM (2022) Remission of obesity and insulin resistance is not sufficient to restore mitochondrial homeostasis in visceral adipose tissue. *Redox Biol* 54:102353. <https://doi.org/10.1016/j.redox.2022.102353>
- Komlodi T, Cardoso LHD, Doerrier C, Moore AL, Rich PR, Gnaiger E (2021) Coupling and pathway control of coenzyme Q redox state and respiration in isolated mitochondria. *Bioenerg Commun* 2021.3. <https://doi.org/10.26124/bec:2021-0003>
- Krumschnabel G, Eigentler A, Fasching M, Gnaiger E (2014) Use of safranin for the assessment of mitochondrial membrane potential by high-resolution respirometry and fluorometry. *Methods Enzymol* 542:163-81. <https://doi.org/10.1016/B978-0-12-416618-9.00009-1>

Copyright ©2026 The author. This is an Open Access preprint (not peer-reviewed) distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. © The author, who has granted BEC preprints an Open Access preprint license in perpetuity.