

A mouse model of acute subdural hematoma under clinically realistic intensive care conditions: insights into effects on mitochondrial respiratory activity

Hannemann W¹, Sharon O², Kress Sa¹, Gröger M¹, Hoffmann A¹, Seifritz A¹, Kranz St¹, Calzia Enrico¹, Radermacher P¹, Kapapa T²

¹Inst Anesthesiol Pathophysiol Process Eng, Ulm Univ, Ulm, Germany; ²Dept Neurosurg, Univ Hosp Ulm, Ulm, Germany. Email: enrico.calzia@uni-ulm.de

Contribution to [MiP2026](#) – Received 2026-09-30 – Online 2026-10-12

Cite: Hannemann W, Sharon O, Kress Sa, Gröger M, Hoffmann A, Seifritz A, Kranz St, Calzia E, Radermacher P, Kapapa T (2026) A mouse model of acute subdural hematoma under clinically realistic intensive care conditions: insights into effects on mitochondrial respiratory activity. BEC preprints 2026.24. <https://doi.org/10.26124/becprep2026-0024>

Introduction Acute subdural hematoma (ASDH) combines primary mass displacement with secondary injury [1], including blood-brain barrier breakdown and excitotoxicity [2] as well as mitochondrial dysfunction [3]. We characterised mitochondrial respiration after ASDH in a mouse model under intensive care monitoring.

Methods ASDH was induced in adult mice ($n = 10$) by subdural injection of autologous blood (10 μ L) using a dual burr-hole model. Intracranial pressure, brain tissue oxygen partial pressure, temperature and arterial pressure were monitored for up to 8 h. Untreated mice ($n = 4$) served as controls. Respiration of homogenised lesion-adjacent and lesion-distant tissue was measured by high-resolution respirometry using a substrate-uncoupler-inhibitor titration (SUIT) protocol [4]. Mass-specific oxygen flux, J_{O_2} , is given in $\text{pmol}\cdot\text{s}^{-1}\cdot\text{mg}^{-1}$ as median [Q1; Q3]. Within-group comparisons were performed by the Wilcoxon signed-rank test ($p < 0.05$).

Results and discussion Preliminary analyses showed markedly lower oxidative phosphorylation (OXPHOS) and electron transfer (ET) capacities in ASDH than in controls (Table 1). Within the ASDH group, respiration was lower in lesion-adjacent than in lesion-distant tissue ($p < 0.05$), whereas controls showed no such difference. The cytochrome c effect, defined as respiration before relative to after cytochrome c addition, was virtually absent in controls but marked in ASDH, particularly in lesion-adjacent tissue ($p < 0.05$). Thus, ASDH markedly impairs mitochondrial respiration with a gradient between lesion-adjacent and lesion-distant tissue, and the cytochrome c response suggests alterations beyond reduced respiratory capacity. This reproducible model enables mechanistic investigations.

Table 1. Respiration of brain tissue homogenates (median [Q1; Q3]). * $p < 0.05$, lesion-adjacent vs lesion-distant (Wilcoxon signed-rank test).

	ASDH, lesion-adjacent	ASDH, lesion-distant	Control, adjacent region	Control, distant region
OXPHOS capacity, J_{O_2} / ($\text{pmol}\cdot\text{s}^{-1}\cdot\text{mg}^{-1}$)	28.2 [19.9; 37.2]*	40.9 [39.9; 44.1]	85.7 [78.5; 91.0]	83.8 [76.9; 88.2]
ET capacity, J_{O_2} / ($\text{pmol}\cdot\text{s}^{-1}\cdot\text{mg}^{-1}$)	32.9 [21.4; 48.7]*	76.3 [60.8; 85.6]	126.2 [120.3; 133.3]	135.3 [127.9; 141.2]
Cytochrome c effect / %	53.1 [46.7; 61.1]*	78.0 [73.8; 84.0]	99.7 [99.0; 100.5]	95.3 [94.8; 96.7]

Keywords acute subdural hematoma; mitochondrial respiration; high-resolution respirometry; traumatic brain injury; mouse model

Author contributions: WH and OS contributed equally. WH and OS conducted the technical aspects of the experiment; WH performed and evaluated the respirometric measurements, and OS performed the neurosurgical procedures. SaK and MG conducted the non-neurosurgical procedures and were responsible for anaesthesiological and intensive care monitoring. AH was responsible for animal welfare and sample collection. AS and StK were responsible for sample processing and data management. EC contributed to experiment planning and supervised the respirometric analyses. PR and TK supervised the entire project as project leaders.

Conflicts of interest The authors declare none.

Funding This work was supported by the German Research Foundation (DFG) Research Unit 5417.

References

- [1] Maas AIR et al (2017) Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol.* 16(12):987-1048 [https://doi.org/10.1016/S1474-4422\(17\)30371-X](https://doi.org/10.1016/S1474-4422(17)30371-X).
- [2] Kapapa T et al (2025) Unravelling secondary brain injury: insights from a human-sized porcine model of acute subdural haematoma. *Cells.* 14(1):17 <https://doi.org/10.3390/cells14010017>.
- [3] Pandya JD, Leung LY, Yang X, Flerlage WJ, Gilsdorf JS, Deng-Bryant Y, Shear DA (2019) Comprehensive profile of acute mitochondrial dysfunction in a preclinical model of severe penetrating TBI. *Front Neurol.* 10:605 <https://doi.org/10.3389/fneur.2019.00605>.
- [4] Buck E, Zügel M, Schumann U, Merz T, Gump AM, Witting A, Steinacker JM, Landwehrmeyer GB, Weydt P, Calzia E, Lindenberg KS (2017) High-resolution respirometry of fine-needle muscle biopsies in pre-manifest Huntington's disease expansion mutation carriers shows normal mitochondrial respiratory function. *PLoS One.* 12(4):e0175248 <https://doi.org/10.1371/journal.pone.0175248>.

Copyright ©2026 The authors. This is an Open Access preprint (not peer-reviewed) distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. © The authors, who have granted BEC preprints an Open Access preprint license in perpetuity